

# Appendix C: Reference Tables

	PAGE
<a href="#">Table of standard atomic weights, 1999</a> . . . . .	3
<a href="#">900.03</a> Temperature corrections for readings of saccharometers (standard at 20°C) . . . . .	6
<a href="#">908.01</a> Alcohol table for calculating percentages of alcohol by volume at 15.56°C (60°F) in mixtures of ethyl alcohol and water from Zeiss immersion refractometer readings and refractive indices at 17.5 –25°C . . . . .	7
<a href="#">909.03</a> Density of carbon dioxide (Parr) . . . . .	13
<a href="#">909.04</a> Correction factors for gasometric determination of carbon dioxide . . . . .	14
<a href="#">913.02</a> Percentages by volume 15.56°C (60°F) of ethyl alcohol corresponding to apparent specific gravity at various temperatures . . . . .	16
<a href="#">921.14</a> Correction table for specific gravity milk (Quevenne lactometer) . . . . .	35
<a href="#">924.10</a> Percentages by weight corresponding to various percentages by volume at 15.56°C (60°F) in mixtures of ethyl alcohol and water . . . . .	36
<a href="#">930.44</a> Total reducing sugar required for complete reduction of 10 mL Soxhlet solution to be used in conjunction with Lane-Eynon general volumetric method . . . . .	37
<a href="#">930.45</a> Total reducing sugar required for complete reduction of 25 mL Soxhlet solution to be used in conjunction with Lane-Eynon general volumetric method . . . . .	38
<a href="#">932.29</a> Jackson-Mathews table of densities of fructose solutions and mean density and expansion coefficients between 20 and 25°C . . . . .	39
<a href="#">933.12</a> Table for determining per cent sucrose in sugar solutions from readings of Zeiss immersion of refractometer at 20°C . . . . .	40
<a href="#">935.70</a> Various strength solutions of the common acids, alkalines, and alcohol . . . . .	41
<a href="#">940.39</a> Hammond table for calculating glucose, fructose, and invert sugar and lactose alone and in the presence of sucrose, with values for maltose from the Munson and Walter table . . . . .	42
<a href="#">942.33</a> Degrees Brix, specific gravity, and degrees Baumé of sugar solutions (Plato table) . . . . .	51
<a href="#">944.16</a> Refractive indices of glucose, fructose, and raffinose hydrate solutions at 20°C . . . . .	57
<a href="#">945.101</a> Table for determining total solids in milk from any given specific gravity and percentage of fat (Shaw and Eckles) . . . . .	58
<a href="#">955.57</a> Optical crystallographic properties of some crystalline drugs . . . . .	60
<a href="#">955.58</a> Table of refractive indices for drugs, arranged according to ascending value of the lowest index. . . . .	65
<a href="#">962.37</a> Density of sucrose solutions at 0 –100°C and 0–70% . . . . .	69
<a href="#">962.38</a> Volume factors for thermal expansion of sucrose solutions up to 100°C . . . . .	70
<a href="#">963.36</a> Refractive indices of invert sugar solutions . . . . .	71

<a href="#">963.37</a>	Nomograph relating absorbance, concentration, and absorptivity (1 cm cell) . . . . .	72
<a href="#">970.90</a>	Specific gravity and degrees Plato of sugar solutions or per cent extract by weight. . . . .	73
<a href="#">990.35A</a>	Concentration of aqueous sucrose solutions (mass fraction, g/100 g) as function of refractive index ( <i>n</i> ) measured at 20°C at wavelength of 589 nm (Na D line) . . . . .	78
<a href="#">990.35B</a>	Concentration of aqueous sucrose solutions (mass fraction, g/100 g) as function of refractive index ( <i>n</i> ) measured at 20°C at wavelength of 546 nm (Hg e line) . . . . .	81
<a href="#">990.36A</a>	Corrections for determination of sucrose in aqueous solutions from refractive index ( <i>n</i> ) readings at wavelength of 589 nm (Na D Line) at temperatures other than 20°* . . . . .	85
<a href="#">990.36B</a>	Corrections for determination of sucrose in aqueous solutions from refractive index ( <i>n</i> ) readings at wavelength of 546 nm (Hg e Line) at temperatures other than 20°* . . . . .	86

**SURPLUS TABLES**  
**(10th Ed.)**

Domke table of apparent specific gravity of sucrose solutions at 20°C (partially replaced by <a href="#">942.33</a> , <a href="#">962.37</a> , and <a href="#">962.38</a> ) . . . . .	<b>43.006</b>
sucrose (0.4 g and 2 g total sugar), lactose, lactose and sucrose (2 mixtures), and maltose (crystalized) Maltose values retained in <a href="#">940.39</a> . . . . .	<b>43.012</b>
Corrections to be subtracted from iodine titer to obtain mg invert sugar by Ofner method . . . . .	<b>43.014</b>
Munson and Walker table for calculating glucose, invert sugar alone, invert sugar in presence of Wein table for determining maltose . . . . .	<b>43.016</b>
Quisumbing and Thomas table for calculating glucose, fructose, invert sugar, lactose, and maltose ( <b>47.024</b> , 11th Ed.). . . . .	<b>43.020</b>
Krober table for determination of pentoses and pentosans . . . . .	<b>43.021</b>
Progressive accumulation of radium emanation . . . . .	<b>43.029</b>
Meissl and Hiller factors for determining invert sugar in materials in which, of total sugars present, more than 1.5% is invert sugar and less than 98.5% is sucrose . . . . .	<b>43.105</b>